


## 14.4 Gradients, Directional Derivatives

Let  $z = f(x, y)$  be given. The partial derivatives  $f_x(a, b)$ ,  $f_y(a, b)$  at some point  $(a, b)$  give the rates of change in the direction of  $x$  and the direction of  $y$ :



$f_x(a, b)$  gives us the rate of change when we walk from  $(a, b)$  in the direction of  $\vec{i}$ ,  $f_y(a, b)$  if we walk in the direction of  $\vec{j}$ .

What is the rate of change if we move from  $(a, b)$  in the direction of some other unit vector  $\vec{u}$ ?

This rate of change is called the directional derivative of  $f(x, y)$  at  $(a, b)$  in the direction of  $\vec{u}$ .


Def: Let  $\vec{u} = u_1\vec{i} + u_2\vec{j}$ ,  $\|\vec{u}\| = 1$ ,  $z = f(x, y)$ ,  $(a, b)$  be given. The directional derivative of  $f(x, y)$  at  $(a, b)$  in the direction of  $\vec{u}$  is:

$$f_{\vec{u}}(a, b) = \lim_{h \rightarrow 0} \frac{f(a + hu_1, b + hu_2) - f(a, b)}{h}$$

$$(a, b) \rightarrow (a + hu_1, b + hu_2)$$

Note: The definition talks about both-sided limit "lim". Thus  $h$  can be positive or negative.

The limit exists if the cross-section of the graph by the vertical plane containing  $\vec{u}$  and  $(a, b, f(a, b))$  has the tangent line at  $(a, b, f(a, b))$ :



Thus  $f_{\vec{u}}(a, b)$  also tells us that if we go in the opposite direction,  $-\vec{u}$ ,  $f(x, y)$  will be changing at the rate  $-f_{\vec{u}}(a, b) = f_{-\vec{u}}(a, b)$

Indeed:

$$\begin{aligned}
 f_{-\vec{u}}(a, b) &= \lim_{h \rightarrow 0} \frac{f(a + h(-u_1), b + h(-u_2)) - f(a, b)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{f(a + (-h)u_1, b + (-h)u_2) - f(a, b)}{-h} \\
 &= -f_{\vec{u}}(a, b)
 \end{aligned}$$

Note, the displacement vector from  $(a, b)$  to  $(a+hu_1, b+hu_2)$  is  $\vec{d} = (hu_1)\vec{i} + (hu_2)\vec{j} = h \cdot \vec{u}$ , so  $\|\vec{d}\| = |h|$  (or more precisely  $|h|$ ).

This is why we need  $\vec{u}$  to be the unit vector so the magnitude of  $h$  is the magnitude of the displacement. Geometrically,  $f_{\vec{u}}(a, b)$  is the slope of the cross-section with the vertical plane parallel to  $\vec{u}$  through  $(a, b, f(a, b))$ . Clearly:

$$f_x(a, b) = f_{\vec{i}}(a, b), \quad f_y(a, b) = f_{\vec{j}}(a, b).$$

Def: Let  $f(x, y)$ ,  $(a, b)$ , and a vector  $\vec{v}$  be given. Then

$$f_{\vec{v}}(a, b) = f \frac{\vec{v}}{\|\vec{v}\|}(a, b).$$

How do we calculate directional derivatives?

Using the so-called gradient vector.

Def: The gradient vector of  $f(x, y)$  at  $(a, b)$  is defined as

$$\text{grad } f(a, b) = f_x(a, b)\vec{i} + f_y(a, b)\vec{j}.$$

-4-

In general:

$$\text{grad } f(x, y) = f_x(x, y) \vec{i} + f_y(x, y) \vec{j}$$

or

$$\text{grad } f = f_x \vec{i} + f_y \vec{j}$$

Ex: Let  $f(x, y) = x^3 y + 3y + x$ . Find  $\text{grad } f(x, y)$ . Find  $\text{grad } f(1, 0)$ .

$$f_x = 3x^2 y + 1, \quad f_y = x^3 + 3$$

$$\text{grad } f = (3x^2 y + 1) \vec{i} + (x^3 + 3) \vec{j}$$

$$\underline{\text{grad } f(1, 0) = \vec{i} + 4\vec{j}}$$

$\text{grad } f$  is a vector!

Th: Let  $f(x, y)$  be differentiable at  $(a, b)$ ,  $\vec{u}$  be the unit vector. Then

$$\begin{aligned} f_{\vec{u}}(a, b) &= \text{grad } f(a, b) \cdot \vec{u} = \\ &= f_x(a, b) u_1 + f_y(a, b) u_2. \end{aligned}$$

Gradient is also denoted:

$$\underline{\text{grad } f = \nabla f}$$

Ex: Let  $\vec{u} = \frac{1}{\sqrt{2}}\vec{i} + \frac{1}{\sqrt{2}}\vec{j}$ ,  $f(x, y) = x^2 + y^2$ .

Find  $f_{\vec{u}}(1, 0)$ .

We use gradient.

$$\text{grad } f(x, y) = 2x\vec{i} + 2y\vec{j}$$

$$\text{grad } f(1, 0) = 2\vec{i}$$


$$f_{\vec{u}}(1, 0) = 2\vec{i} \cdot \vec{u} = 2 \cdot \frac{1}{\sqrt{2}} = \sqrt{2}.$$

So 'finding' directional derivatives is as easy as finding gradients.

Let  $f(x, y)$ , differentiable, be given. Then we have the gradient vector at each point:


$$\nabla f(x, y)$$

So we have a vector field - the gradient field on the  $xy$ -plane:



## Gradient Geometrically

Let's have  $f(x, y)$ ,  $(a, b)$ ,  $\nabla f(a, b)$ :



How about directional derivatives at  $(a, b)$ ?

Let  $\vec{u}$  be a unit vector.

$$f_{\vec{u}}(a, b) = \nabla f(a, b) \cdot \vec{u} = \|\nabla f(a, b)\| \cdot \|\vec{u}\| \cos(\theta)$$

$f_{\vec{u}}(a, b)$  is the largest if  $\cos(\theta) = 1$ ; that is,

when  $\theta = 0$ , so  $\vec{u}$  is parallel to  $\nabla f(a, b)$ .

Thus:


Th: Let  $f(x, y)$  be differentiable at  $(a, b)$ ,  $\nabla f(a, b) \neq \vec{0}$ .

Then:

- (1)  $\nabla f(a, b)$  points in the direction of the largest rate of change of  $f$  at  $(a, b)$ . (The direction of the fastest growth.)
- (2) This largest rate of change is  $\|\nabla f(a, b)\|$ .
- (3)  $\nabla f(a, b)$  is perpendicular to the contour line of  $f(x, y)$  through  $(a, b)$ .

~~7~~

Why (3)? Since  $f(x,y)$  is differentiable at  $(a,b)$ , locally near  $(a,b)$  the graph of  $z=f(x,y)$  is flat. Thus, locally contour lines are parallel:





You obtain fastest growth by moving perpendicularly to contours in the direction of increasing values. So  $\nabla f(a,b)$  that points toward fastest increase must be perpendicular to contours.

that points toward fastest increase must be perpendicular to contours.

Ex: Let  $f(x,y) = x^2 + y^2$ . Sketch the contour diagram and the gradient field in one coordinate system.

$$\nabla f(x,y) = 2x\vec{i} + 2y\vec{j}$$

parallel the vector from  $(0,0)$  to  $(x,y)$ .



Magnitudes of vectors are scaled.

Ex A square metal plate is placed in the  $xy$ -plane in such a way that  $0 \leq x \leq 3$ ,  $0 \leq y \leq 3$ . (All measurements in meters.) The temperature at each point  $(x, y)$  of the plane is given by:

$$T(x, y) = \frac{100}{x^2 + y^2 + 1} \text{ in } ^\circ\text{F.}$$


- (a) Find the direction of the greatest increase in temp. at  $(1, 2)$ .  
What is the greatest rate of increase at the point  $(1, 2)$ ?
- (b) Find the rate of increase at  $(1, 2)$  in the direction  $\vec{u} = \frac{2}{\sqrt{5}}\vec{i} + \frac{1}{\sqrt{5}}\vec{j}$ .

Clearly, we have to find the gradient  $\nabla T(x, y)$  first.

$$\begin{aligned} T_x(x, y) &= \frac{\partial}{\partial x} \left[ \frac{100}{x^2 + y^2 + 1} \right] = \frac{\partial}{\partial x} \left[ 100(x^2 + y^2 + 1)^{-1} \right] = \\ &= 100 \cdot -(x^2 + y^2 + 1)^{-2} \cdot 2x = \\ &= -\frac{100}{(x^2 + y^2 + 1)^2} \cdot 2x = T_x(x, y) \end{aligned}$$



$$T_y(x, y) = -\frac{100}{(x^2 + y^2 + 1)^2} \cdot 2y$$

$$\nabla T(x, y) = -\frac{100}{(x^2 + y^2 + 1)^2} (2x\vec{i} + 2y\vec{j})$$






-9-



Contours are concentric circles centered at (0,0).  
At each point the gradient  $\nabla T(x, y)$  points toward the origin.



$$\nabla T(1,2) = -\frac{200}{36} \vec{i} - \frac{400}{36} \vec{j}$$

The direction of the greatest increase in temperature at (1,2). (Toward the origin). This answers (a).

$$(b) \quad \|\nabla T(1,2)\| = \sqrt{\left(\frac{200}{36}\right)^2 + \left(\frac{400}{36}\right)^2} \approx 12.42 \frac{^\circ\text{F}}{\text{m}}$$

$$(c) \quad T_{\vec{u}}(1,2) = \nabla T(1,2) \cdot \vec{u} =$$

$$= -\frac{200}{36} \cdot \frac{2}{\sqrt{5}} - \frac{400}{36} \cdot \frac{1}{\sqrt{5}} \approx -9.93 \frac{^\circ\text{F}}{\text{m}}$$


## 14.5 Gradients of Functions of Three Variables

Ex:  $f(x, y, z) = x^2 + y^2 + z^2$

We cannot "graph" functions of three variables.

We can graph their level surface for a given  $k$ :

$$x^2 + y^2 + z^2 = k$$



We can, of course, define three partial derivatives at each point  $(a, b, c)$ :

$$f_x(a, b, c), f_y(a, b, c), f_z(a, b, c)$$

by fixing two variables and differentiating with respect to the third. For example:

$$f_z(a, b, c) = \left. \frac{d}{dz} [f(a, b, z)] \right|_{z=c}$$

In our example:

$$f_x(x, y, z) = 2x, f_y(x, y, z) = 2y, f_z(x, y, z) = 2z.$$

We can define the directional derivative of

a function of three variables  $f(x, y, z)$  in the direction of  $\vec{u} = u_1 \vec{i} + u_2 \vec{j} + u_3 \vec{k}$ ,

$$\|\vec{u}\| = 1:$$

$$f_{\vec{u}}(a, b, c) = \lim_{h \rightarrow 0} \frac{f(a+hu_1, b+hu_2, c+hu_3) - f(a, b, c)}{h}$$

This is the rate of change in the direction of  $\vec{u}$ .

We can define the gradient vector:

$$\nabla f(x, y, z) = f_x(x, y, z)\vec{i} + f_y(x, y, z)\vec{j} + f_z(x, y, z)\vec{k}.$$

In our example,  $f(x, y, z) = x^2 + y^2 + z^2$ ,

$$\nabla f = 2x\vec{i} + 2y\vec{j} + 2z\vec{k}.$$

As before, if  $f(x, y, z)$  is differentiable at  $(a, b, c)$ :

$$f_{\vec{u}}(a, b, c) = \nabla f(a, b, c) \cdot \vec{u} \quad \frac{\text{units of } f}{\text{units of dist in dir } \vec{u}}$$

As before:

If  $\nabla f(a, b, c) \neq \vec{0}$ , then

- $\nabla f(a, b, c)$  points in the direction of the greatest rate of change of  $f$  at  $(a, b, c)$ .
- $\|\nabla f(a, b, c)\|$  is this greatest rate of change
- $\nabla f(a, b, c)$  is perpendicular to the level surface through  $(a, b, c)$ .


Ex Suppose that the function  $F(x, y, z) = x^2 + y^4 + x^2 z^2$  gives concentration of salt, in gr/gal, at any point  $(x, y, z)$  of a rectangular tank of water occupying the region

$$-2 \leq x \leq 2, \quad -2 \leq y \leq 2, \quad 0 \leq z \leq 2.$$

(All measurements in meters.) Suppose you are at the point  $(-1, 1, 1)$ .

(a) In what direction should you move if you want the concentration to increase the fastest?

(b) If you move from  $(-1, 1, 1)$  toward the origin  $(0, 0, 0)$ , how fast is the concentration changing?



(a)

$$\begin{aligned} \nabla F(x, y, z) &= \\ &= (2x + 2xz^2)\vec{i} + 4y^3\vec{j} + 2x^2z\vec{k} \end{aligned}$$

$$\nabla F(-1, 1, 1) = -4\vec{i} + 4\vec{j} + 2\vec{k}$$

↑ The direction of greatest increase in concentration.

$$\|\nabla F(-1, 1, 1)\| = \sqrt{16 + 16 + 4} = 6 \frac{\text{g/gal}}{\text{m}}$$

$$(b) \vec{v} = \overrightarrow{(-1, 1, 1)(0, 0, 0)} = \vec{i} - \vec{j} - \vec{k}, \quad \|\vec{v}\| = \sqrt{3}$$

$$\vec{u} = \frac{1}{\sqrt{3}} \vec{i} - \frac{1}{\sqrt{3}} \vec{j} - \frac{1}{\sqrt{3}} \vec{k}$$

$$F_{\vec{u}}(-1, 1, 1) = F_{\vec{u}}(-1, 1, 1) = \nabla F(-1, 1, 1) \cdot \vec{u} =$$

$$= -\frac{4}{\sqrt{3}} - \frac{4}{\sqrt{3}} - \frac{2}{\sqrt{3}} = -\frac{10}{\sqrt{3}} \approx -5.77 \frac{\text{g/gal}}{\text{m}}$$

Ex: Find the equation of the tangent plane to the ellipsoid  $x^2 + 2y^2 + z^2 = 15$  at  $(2, 1, 3)$ .

The ellipsoid is the level surface of


$$F(x, y, z) = x^2 + 2y^2 + z^2$$

$$\vec{n} = \nabla F(2, 1, 3)$$

$$\nabla F = 2x\vec{i} + 4y\vec{j} + 2z\vec{k}$$

$$\vec{n} = \nabla F(2, 1, 3) = 4\vec{i} + 4\vec{j} + 6\vec{k}$$

$$P = (2, 1, 3)$$



$$\text{Plane: } 4(x-2) + 4(y-1) + 6(z-3) = 0.$$


---