

§ 13.5 The Exponential Function

Idea: Extend the usual exponential f s e^x from calculus to the complex numbers.

For $z = x + iy \in \mathbb{C}$, define the exponential of z , e^z or $\exp z$ by

$$e^z := e^x (\cos y + i \sin y).$$

Note that if $z = x \in \mathbb{R}$, then $y = 0$

so $\cos y = 1$, $\sin y = 0$ and

$$e^z = e^x,$$

so our new definition does indeed extend e^x .

We can define the derivative of a complex fn by analogy with the derivative from calculus.

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \quad (h \text{ is complex}).$$

With this defn one can show e^z is differentiable everywhere and

$$\frac{d}{dz} (e^z) = e^z, \quad z \in \mathbb{C}$$

just like $\frac{d}{dx} (e^x) = e^x, \quad x \in \mathbb{R}.$

Fns which are differentiable in this sense are often called analytic or holomorphic.

Fns like e^z which are analytic everywhere are called entire fns.

If $z_1 = x_1 + iy_1$, $z_2 = x_2 + iy_2$, then

$$e^{z_1} e^{z_2} = e^{x_1 + iy_1} e^{x_2 + iy_2}$$

$$= e^{x_1} (\cos y_1 + i \sin y_1) \cdot e^{x_2} (\cos y_2 + i \sin y_2)$$

$$= e^{x_1} e^{x_2} (\cos y_1 \cos y_2 - \sin y_1 \sin y_2 + i (\cos y_1 \sin y_2 + \sin y_1 \cos y_2))$$

$$= e^{x_1 + x_2} (\cos(y_1 + y_2) + i \sin(y_1 + y_2))$$

by compound angle & $e^a e^b = e^{a+b}$

$$= e^{x_1 + x_2 + i(y_1 + y_2)}$$

$$= e^{z_1 + z_2}$$

So $e^{z_1 + z_2} = e^{z_1} e^{z_2}$ (like $e^{a+b} = e^a e^b$ in calculus)

Note if $z = x + iy$, then

$$e^z = e^x e^{iy}$$

If $z = iy$, $x = 0$ and

$$e^z = \cos y + i \sin y, \text{ so}$$

$$e^{iy} = \cos y + i \sin y$$

Hence for a cplx no $z = r(\cos \theta + i \sin \theta)$
in polar form, we may write

$$z = r e^{i\theta}$$

Special cases of the above are

$$e^{i\pi} = -1,$$

Euler's formula

$$e^{2\pi i} = 1, \quad e^{i\frac{\pi}{2}} = i, \quad e^{-i\frac{\pi}{2}} = -i$$

Note that since

$$e^z = e^x (\cos y + i \sin y)$$

is already in polar form, we see that

$$|e^z| = e^x, \quad \arg(e^z) = y \quad \text{up to integer multiples of } 2\pi$$

In particular, $|e^{iy}| = 1$ ($x = 0$).


Also

$$\begin{aligned} e^{z+2\pi i} &= e^x (\cos(y+2\pi) + i \sin(y+2\pi)) \\ &= e^x (\cos y + i \sin y) \quad \text{by periodicity of } \cos, \sin \\ &= e^z. \end{aligned}$$

So $e^{z+2\pi i} = e^z$. Say e^z is periodic with period 2π .

Thus e^z will assume all its possible values on a horizontal strip of width 2π , e.g. the strip

$$S = \{ z = x + iy : -\pi < y \leq \pi \}.$$



Finally note that since $|e^z| = e^x$ and $e^x \neq 0$ for all $x \in \mathbb{R}$, e^z is never 0 in value. Say e^z is entire and non-vanishing.