

7.2 Point Estimates for μ and σ^2 .

The most important function in all of statistics:

$$f(y) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(y-\mu)^2}$$

Recall the MLE estimates for a given random sample of size n :

$$\hat{\mu} = \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$

and

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$$

Recall that $\hat{\sigma}^2$ is biased. In practice, the sample variance S^2 is used

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$$

171

Fitting a Normal Distribution to Data

GIVEN: n data points y_1, \dots, y_n

GROUP data points into k equal-width classes

SET:

m_i = midpoint of i -th class

f_i = number of observations in i -th class.

COMPUTE the density for each class

$$\text{class density} = \frac{\text{class frequency}}{\text{class width} \times n}$$

or

$$d_i = \frac{f_i}{\Delta \times n}$$

173

To estimate μ use the grouped sample mean

$$\bar{y}_g = \frac{1}{n} \sum_{i=1}^k f_i m_i$$

To estimate σ^2 use the grouped sample variance

$$\begin{aligned} s_g^2 &= \frac{1}{n-1} \sum_{i=1}^k f_i (m_i - \bar{y}_g)^2 \\ &= \frac{1}{n(n-1)} \left(n \sum_{i=1}^k f_i m_i^2 - \left(\sum_{i=1}^k f_i m_i \right)^2 \right) \end{aligned}$$

In the previous example,

$$\bar{y}_g = 67.60 \quad \text{and} \quad s_g^2 = 6.656$$

175

Theorem 7.2.1 Let Y_1, Y_2, \dots, Y_n be a random sample from a normal distribution with mean μ and variance σ^2 . Then \bar{Y} , the MLE for μ , is unbiased, efficient, and consistent. If in addition σ^2 is known, then \bar{Y} is sufficient.

Here we prove consistency, that is, $\lim_{n \rightarrow \infty} P(|\bar{Y}_n - \mu|) = 1$. Chebyshev's inequality gives

$$P(|\bar{Y}_n - \mu| < \epsilon) \geq 1 - \frac{\text{Var}(\bar{Y})}{\epsilon^2}$$

Since $\text{Var}(\bar{Y}) = \sigma^2/n$, we have

$$1 - \frac{\sigma^2}{n\epsilon^2} \leq P(|\bar{Y}_n - \mu| < \epsilon) \leq 1$$

Now take limit on all three terms. The inequalities are preserved, and the outside limits give 1. By the "squeeze theorem" the middle limit is also 1.

172




FIGURE 2.5.8




FIGURE 2.5.9

174

Height of US Army recruits. Is it normally distributed?

TABLE 7.2.1

Height (in.)	Midpoint, m_i	Frequency, f_i	Density
54.0 ≤ y < 55.0	54.5	5	0.00019
55.0 ≤ y < 56.0	55.5	3	0.00012
56.0 ≤ y < 57.0	56.5	7	0.00027
57.0 ≤ y < 58.0	57.5	6	0.00023
58.0 ≤ y < 59.0	58.5	10	0.00039
59.0 ≤ y < 60.0	59.5	15	0.00058
60.0 ≤ y < 61.0	60.5	50	0.00193
61.0 ≤ y < 62.0	61.5	57	0.00225
62.0 ≤ y < 63.0	62.5	1237	0.04859
63.0 ≤ y < 64.0	63.5	1047	0.07524
64.0 ≤ y < 65.0	64.5	3019	0.11666
65.0 ≤ y < 66.0	65.5	3475	0.13428
66.0 ≤ y < 67.0	66.5	4054	0.15666
67.0 ≤ y < 68.0	67.5	9631	0.31031
68.0 ≤ y < 69.0	68.5	3133	0.12107
69.0 ≤ y < 70.0	69.5	2075	0.08018
70.0 ≤ y < 71.0	70.5	1485	0.05738
71.0 ≤ y < 72.0	71.5	680	0.02628
72.0 ≤ y < 73.0	72.5	343	0.01325
73.0 ≤ y < 74.0	73.5	118	0.00456
74.0 ≤ y < 75.0	74.5	42	0.00162
75.0 ≤ y < 76.0	75.5	9	0.00035
76.0 ≤ y < 77.0	76.5	6	0.00023
77.0 ≤ y < 78.0	77.5	2	0.00008
		25878	

176




FIGURE 7.2.1

177

178

Case Study 7.2.2: $\frac{\bar{Y} - \mu}{\sigma/\sqrt{n}}$ vs. $\frac{\bar{Y} - \mu}{S/\sqrt{n}}$




FIGURE 7.2.2

179