

17.1-17.2 Parametrized Curves, Motion


You are familiar with parametric representations of curves on the xy -plane.

Ex: What curve on the xy -plane is described by:

$$x = \cos t, \quad y = \sin t, \quad t \text{ in } [0, 2\pi]$$

↑
the parameter

Since $\cos^2 t + \sin^2 t = 1$, each point on the parametric curve is on the unit circle:




$$t=0 \rightarrow (1, 0)$$

As t increases, we move counter-clockwise; at $t=2\pi$, we are back at $(1, 0)$.

Of course, the unit circle has other parametric representations.

For example:

$$x = \sin t, \quad y = \cos t, \quad t \in [0, 2\pi]$$




Same path, different parametrization.

Parametrizations provide a very convenient way of representing curves in the xyz -space.

A parametric curve in the xyz -space is a curve described by parametric equations:

$$x = f(t), \quad y = g(t), \quad z = h(t)$$

where the parameter t changes in an interval I .




For each t we have a point $(f(t), g(t), h(t))$ on the curve C . As t changes, the point moves along C describing a motion along the path C .

Ex: What curve is described by

$$x = \cos t, \quad y = \sin t, \quad z = t, \quad t \geq 0.$$

What motion along the curve is described by the parametrization?



Note that the projection onto the xy -plane:

$$x = \cos t, \quad y = \sin t$$

moves ccw about the unit circle. At the same time z increases so we move up.

A helix in the xyz -space.

The same helix can be parametrized by;

Ex: $x = \cos(2t), y = \sin(2t), z = 2t, t \geq 0.$

In the latter parametrization we traverse the helix twice as fast.

Parametrization in Vector Form


Let a parametrized curve

$$x = f(t), y = g(t), z = h(t), t \in I$$

be given. We can write the parametrization

as: $\vec{r}(t) = f(t)\vec{i} + g(t)\vec{j} + h(t)\vec{k}, t \in I.$

$\vec{r}(t)$ is called the position vector:



The position vector traces the curve as t changes.

The simplest way to describe a straight line in 3D is by a parametric representation.

Let L be the line passing through a point (x_0, y_0, z_0) and parallel to $\vec{w} = w_1\vec{i} + w_2\vec{j} + w_3\vec{k}$. Then

$$L : x(t) = x_0 + tw_1, \quad y(t) = y_0 + tw_2, \quad z(t) = z_0 + tw_3,$$

$$-\infty < t < +\infty.$$


In the vector form:

$$L: \vec{r}(t) = \vec{r}_0 + t\vec{w}, \quad \vec{r}_0 = x_0\vec{i} + y_0\vec{j} + z_0\vec{k},$$

$$-\infty < t < +\infty$$

Of course, the displacement vector from (x_0, y_0, z_0) to $(x(t), y(t), z(t))$ is $tw_1\vec{i} + tw_2\vec{j} + tw_3\vec{k} \parallel \vec{w}$. So

we start from (x_0, y_0, z_0) and move t units in the direction of \vec{w} .



\vec{r}_0 is the position vector of the initial point.

Ex: Let $P_0 = (2, -1, 3)$, $P_1 = (-1, 5, 4)$. Find

a parametric representation of:

(a) The line, L , through P_0, P_1

(b) The segment, S , from P_0 to P_1 .

(a) $\vec{w} = \overrightarrow{(2, -1, 3) - (-1, 5, 4)} = -3\vec{i} + 6\vec{j} + \vec{k}$ - the direction vector

$P_0 = (x_0, y_0, z_0) = (2, -1, 3)$ - initial point

$\vec{r}_0 = 2\vec{i} - \vec{j} + 3\vec{k}$ - the position vector of the initial point.

Parametric representation:

$L: \vec{r}(t) = \vec{r}_0 + t\vec{w}, \quad -\infty < t < +\infty$

Thus:

$L: \vec{r}(t) = 2\vec{i} - \vec{j} + 3\vec{k} + t(-3)\vec{i} + t6\vec{j} + t\vec{k}$

$L: \vec{r}(t) = (2-3t)\vec{i} + (-1+6t)\vec{j} + (3+t)\vec{k}$

In non-vector form:

$L: x(t) = 2-3t, y(t) = -1+6t, z(t) = 3+t$
 $-\infty < t < +\infty$

(b) Since \vec{w} is $\overrightarrow{P_0P_1}$, in the representation


$\vec{r}(t) = \vec{r}_0 + t\vec{w}$

we are at P_0 for $t=0$ and at P_1 at $t=1$.

So:


$S: \vec{r}(t) = (2-3t)\vec{i} + (-1+6t)\vec{j} + (3+t)\vec{k}$

$0 \leq t \leq 1.$



Note: A parametric representation gives us not only a path (L or S) but also a specific motion along the path.

Ex: Find a parametric equation of the circle of radius 3 parallel to the xy -plane centered at $(0, 0, 2)$.



$$x = 3 \cos t, \quad y = 3 \sin t, \quad z = 2$$

$$0 \leq t \leq 2\pi$$

Or

$$\vec{r}(t) = (3 \cos t) \vec{i} + (3 \sin t) \vec{j} + 2 \vec{k}$$

$$0 \leq t \leq 2\pi,$$

Of course, we use parametric representations to describe motion in xyz -space. The parameter t denotes time. What is the velocity and the acceleration?

Def: Let $\vec{r}(t) = f(t)\vec{i} + g(t)\vec{j} + h(t)\vec{k}$ be the position vector of an object at time t . The velocity vector, $\vec{v}(t)$, at time t is defined as:

$$\vec{v}(t) = \lim_{\Delta t \rightarrow 0} \frac{\vec{r}(t + \Delta t) - \vec{r}(t)}{\Delta t} = \vec{r}'(t) = \frac{d\vec{r}}{dt}.$$


In terms of coordinates;

$$\vec{v}(t) = f'(t)\vec{i} + g'(t)\vec{j} + h'(t)\vec{k} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}.$$

Speed is defined as:

$$\text{Speed} = \|\vec{v}(t)\| = \sqrt{f'(t)^2 + g'(t)^2 + h'(t)^2}.$$

$\vec{v}(t)$ is tangent to the path at each point;



Becomes more and more

tangent as $\Delta t \rightarrow 0$.

$\vec{v}(t)$ points in the direction of the motion.

Def: Let $\vec{r}(t) = f(t)\vec{i} + g(t)\vec{j} + h(t)\vec{k}$ be the position vector of an object at time t . The acceleration vector $\vec{a}(t)$ is defined as

$$\vec{a}(t) = \lim_{\Delta t \rightarrow 0} \frac{\vec{v}(t+\Delta t) - \vec{v}(t)}{\Delta t} = \frac{d\vec{v}}{dt} = \vec{v}'(t) = \vec{r}''(t) = \frac{d^2\vec{r}}{dt^2}$$


In terms of coordinates:

$$\vec{a}(t) = f''(t)\vec{i} + g''(t)\vec{j} + h''(t)\vec{k}$$

$\vec{a}(t)$ reflects the changes in both direction and magnitude of the velocity $\vec{v}(t)$.

Ex A particle starts at the point $P = (3, 2, -5)$ and moves along a straight line toward $Q = (5, 7, -2)$ at a speed of $5 \frac{\text{cm}}{\text{sec}}$. Let x, y, z be measured in cm.

- (a) Find the particle's velocity vector.
- (b) Find parametric equations for the particle's motion.



$$\vec{v}(t) \parallel \vec{PQ} = 2\vec{i} + 5\vec{j} + 3\vec{k}$$

t in seconds.

$$\|\vec{v}(t)\| = 5 \frac{\text{cm}}{\text{sec}}$$

$$\vec{v}(t) \equiv \vec{v} = c2\vec{i} + c5\vec{j} + c3\vec{k} \quad c = ?$$

- 8 -

$$\|\vec{v}\| = \sqrt{4c^2 + 25c^2 + 9c^2} = c\sqrt{38} = 5 \rightarrow c = \frac{5}{\sqrt{38}}$$

$$\vec{v}(t) \equiv \frac{5}{\sqrt{38}} \cdot \vec{PQ} = \frac{10}{\sqrt{38}} \vec{i} + \frac{25}{\sqrt{38}} \vec{j} + \frac{15}{\sqrt{38}} \vec{k}$$



$$\vec{a}(t) = \vec{0}$$

Indeed, neither the direction nor the magnitude of the velocity changes.

Ex: Let $\vec{r}(t) = (\cos t)\vec{i} + (\sin t)\vec{j}$. Find $\vec{v}(t)$, $\vec{a}(t)$,
 $t \in [0, 2\pi]$

$$x = \cos t, \quad y = \sin t, \quad z = 0$$

$$\vec{v}(t) = -\sin(t)\vec{i} + \cos(t)\vec{j}$$



$$\vec{a}(t) = -\cos(t)\vec{i} - \sin(t)\vec{j} \quad \parallel \vec{r}(t)$$

$$\vec{a}(t) \perp \vec{v}(t)$$


$\|\vec{v}(t)\| \equiv 1$. $\vec{a}(t)$ reflects changes in the direction of $\vec{v}(t)$.

$$\text{Length of } C = \int_0^{2\pi} \|\vec{v}(t)\| dt = \int_0^{2\pi} (\sin^2(t) + \cos^2(t)) dt = 2\pi.$$

Ex: Consider the motion of an object:

$$\vec{r}(t) = t^2 \vec{j}, \quad t \geq 0$$

Find $\vec{v}(t)$ and $\vec{a}(t)$. Describe the motion.



The object moves faster and faster along the y axis.

$$\vec{v}(t) = 2t \vec{j} \quad - \text{parallel to the } y\text{-axis}$$

$$s(t) = \sqrt{4t^2} = 2t \quad - \text{speed increasing}$$

$$\vec{a}(t) = 2 \vec{j} \quad - \text{acceleration not } \vec{0} \text{ as the magnitude of } \vec{v}(t) \text{ changes.}$$

$\vec{a}(t)$ reflects changes in the magnitude of $\vec{v}(t)$.

Let $\vec{v}(t)$ be the velocity in a motion. Then

$$\text{Distance traveled between } t=a \text{ and } t=b = \int_a^b \|\vec{v}(t)\| dt$$

Suppose an object moves along a curve C , covers the curve once for t in $[a, b]$, then the length of C

$$\text{Length of } C = \int_a^b \|\vec{v}(t)\| dt$$
